

Annual Report of the Great Lakes Regional Water Use Database

Representing 2014 Water Use Data

In Liters

Prepared by the Great Lakes Commission for the Great Lakes-St. Lawrence River Water Resources Regional Body and the Great Lakes-St. Lawrence River Basin Water Resources Council



Table of Contents

| Acknowledgments | 3 |
|---|------|
| Preface | 3 |
| Overview | 4 |
| Improving Data Quality | 4 |
| Data Reporting by Jurisdiction | 5 |
| Great Lakes Regional Water Use for 2014 | 6 |
| Lake Watershed Summaries | 11 |
| Lake Superior | . 11 |
| Lake Michigan | 13 |
| Lake Huron | . 15 |
| Lake Erie | . 17 |
| Lake Ontario | . 19 |
| St. Lawrence River | 21 |
| Jurisdiction Reports | 23 |
| Illinois | 23 |
| Indiana | 24 |
| Michigan | 26 |
| Minnesota | 28 |
| New York | 29 |
| Ohio | . 30 |
| Ontario | . 31 |
| Pennsylvania | 33 |
| Québec | . 34 |
| Wisconsin | . 35 |
| Appendices | 36 |

Acknowledgments

The Great Lakes Commission thanks the states and provinces for their commitment in collecting and reporting the water use data included in this report. Appreciation is extended to Jim Casey and Dan Injerd (Illinois), Allison Mann (Indiana), Andrew LeBaron (Michigan), Sean Hunt (Minnesota), Rich Kruzansky (New York), Lenn Black (Ohio), Jonathan Staples (Ontario), Dave Jostenski and Michael Hill (Pennsylvania), Marc-Olivier Bédard (Québec), Kathleen Mooney and Shaili Pfeiffer (Wisconsin) for their time and effort in preparing and reviewing the 2014 water use data. Special thanks go out to Carol Luukkonen and Kim Shaffer at the U.S. Geological Survey who conducted an independent review of the 2014 metadata and this 2014 annual water use report. Special thanks go to the Conference of Great Lakes-St. Lawrence Governors and Premiers for their financial support of this effort and careful review of this report.

Preface

This is the Annual Report of the Great Lakes-St. Lawrence River Regional Water Use Database, representing 2014 water use data. These data are provided by the Great Lakes-St. Lawrence River states and provinces to the Great Lakes Commission (GLC), which serves as the database repository, under the Great Lakes-St. Lawrence River Basin Water Resources Compact (Compact) and the Great Lakes St. Lawrence River Basin Sustainable Water Resources Agreement (Agreement).

The Great Lakes-St. Lawrence River Regional Water Use Database has been operational since 1988. It was established by the states and provinces in response to a provision of the 1985 Great Lakes Charter, which called for the establishment and maintenance of a regional system for the collection of data on major water uses, diversions and consumptive uses in the binational Great Lakes-St. Lawrence River Basin (Basin). The Charter (a precursor to the Compact and Agreement) was a non-binding, "good faith" agreement signed by the Great Lakes governors and premiers that set forth a series of principles and procedures for strengthening water management activities in the Basin. The Charter envisioned a centralized database as an important tool to support a regional water resources management program that guides the future development, management and conservation of the water resources of the Basin. In 1988, the GLC was selected to serve as the repository for the regional water use database. The maintenance and operation of the database has been provided by the GLC since that time, in partial fulfillment of the Charter obligations and since 2009 supporting the requirements under the Compact and Agreement through a new set of water use data collection and reporting protocols.

After two decades of collecting water use data and issuing the annual water use reports under the Charter, the database has been revised and upgraded to meet the requirements set forth by the Compact and Agreement. In 2008, to help implement the needed improvements in jurisdictional water use data collection and reporting programs, the Council of Great Lakes Governors, through its Great Lakes Water Use Information Initiative, led the states and provinces through a process that culminated in the drafting of new water use data collection and reporting protocols. The Compact Council and Regional Body adopted the new protocols in 2009. The protocols offer guidance to ensure that water use data provided to the database repository by the states and provinces is accurate, of the highest quality, and reported in a common and consistent manner. The 2014 annual water use report presents the third dataset that was assembled using the 2009 water use data collection and reporting protocols.

While the common data protocols are an important step in support of a more robust regional water management regime, it is recognized that much additional work needs to be done and that improvements in data collection, reporting, quality, accuracy and compatibility must continue to occur. Additional information describing the improvements to the data collected under the 2009 regional water use data collection and reporting protocols is provided in Appendix A. The following section describes the progress made in 2015 to improve data quality and describes the quality of the data for the 2014 annual report.

Overview

Improving Data Quality

Together with the Conference of Great Lakes-St. Lawrence Governors and Premiers, the GLC is working with the Great Lakes and St. Lawrence River states and provinces to improve data collection, reporting, quality, accuracy and compatibility. To guide the preparation of 2014 data and this report, several steps have been made to improve data quality.

An important first step to improve data quality was the development of a process to collect metadata for the 2014 water use data. For purposes of this report, metadata is information that describes water use data and includes information related to data sources, reporting compliance rates by water use sector, documenting the year for which the data is collected, any significant changes in the data between the current year and previous years, and describing reasons for those changes. To achieve this purpose, the GLC created an online data management system that assists in the documentation of metadata. For this report, the states and provinces have submitted metadata along with the associated 2014 water use data to the GLC. Through the creation of metadata, states, provinces and the GLC were better able to identify and correct errors in 2013 and 2014 datasets.

Another step to improve data quality was to work with water use specialists at the U.S. Geological Survey to conduct a review of the 2014 metadata. Based upon this review, several of the jurisdictions updated their datasets or modified their metadata. This review has also provided some recommendations to improve the documentation of the metadata which will be discussed further with the states and provinces.

The GLC will continue to work with the states and province to identify additional areas for improvement and will carry out the activities begun in 2014. While this report contains the best available information as of its publishing date (December, 2015), the states and provinces may continue to update their 2014 data housed in the online database. Therefore, discrepancies between the data online and those summarized in this report may appear. Data summarized in this report are available to download from the GLC website at projects.glc.org/waterusedata.

Data Reporting by Jurisdiction

The states and provinces have water use reporting programs in place that requires water users to report their water use each year to their jurisdiction. The reporting compliance, i.e., the percentage of water users submitting the water use reports to the jurisdiction, varies across the Basin and impacts the quality of the data. Reporting compliance varies across sectors for most jurisdictions. Illinois, Minnesota and Ohio report a 100 percent reporting compliance for all sectors. Since Québec is still ramping up its reporting program, reporting compliance could not be calculated for the commercial and institutional, livestock and other self-supply sectors.

The actual year in which the data was collected impacts the comparability of data across the Basin. Except for Ontario and Michigan, all jurisdictions are reporting 2014 data for this annual report. The tables 1 and 2 summarize reporting compliance rates and the actual year the data was collected by jurisdiction.

| Jurisdiction | Actual Data Year | Reporting Compliance |
|--------------|--|------------------------|
| Illinois | 2014 for all sectors | 100% for all sectors |
| Indiana | 2014 for all sectors | Varies across sectors* |
| Michigan | 2014 for most sectors 2012-2013 for self-supply livestock, aquaculture facilities | Varies across sectors* |
| Minnesota | 2014 for all sectors | 100% for all sectors |
| New York | 2014 for all sectors | Varies across sectors* |
| Ohio | 2014 for all sectors | 100% for all sectors |
| Ontario | 2013 for most sectors 2009 data used for public supply intrabasin diversions | Varies across sectors* |
| Pennsylvania | 2014 for all sectors | Varies across sectors* |
| Quebec | 2014 for all sectors | Varies across sectors* |
| Wisconsin | 2014 for all sectors | Varies across sectors* |

Table 1. Actual Year of Water Use Data by Jurisdiction


*See Table 2 for reporting compliance to the jurisdiction by specific water use sector

Table 2. Reporting Compliance to the Jurisdiction by Water Use Sector

| Sector | IN | MI | NY | ONT | PA | WI | QC |
|---|-----|----|-----|-----|-----|-----|-------|
| Public Water Supply | 97 | 99 | 89 | 98 | 94 | 100 | 100 |
| Self-Supply Commercial & Institutional | 90 | 95 | 93 | 100 | 100 | 96 | N/A** |
| Self-Supply Irrigation | 95 | 85 | 56 | 99 | 21 | 96 | N/A |
| Self-Supply Livestock | 75 | 40 | 76 | 99 | 82 | 97 | - |
| Self-Supply Industrial | 95 | 95 | 92 | 88 | 91 | 94 | 100 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 100 | 99 | 92 | 100 | - | 100 | - |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 100 | 99 | 92 | - | - | 100 | - |
| Off-Stream Hydroelectric Power Production | _* | - | 100 | - | - | - | - |
| In-Stream Hydroelectric Water Use | - | - | 100 | 95 | - | - | - |
| Other Self-Supply | 81 | 95 | 62 | 94 | - | 98 | N/A |

*A blank indicates that the jurisdiction did not report any water use figures for that particular sector.

******N/A indicated that reporting compliance percentage could not be calculated.



Great Lakes Regional Water Use for 2014


Figure 1. Great Lakes-St. Lawrence River Basin

The Great Lakes and the St. Lawrence River – the world's largest freshwater system – span an area of 750,000 square kilometers (289,600 square miles). Its total volume is 25 quadrillion liters, an amount that would fill 9 billion Olympic size swimming pools.¹

In 2014, the total reported withdrawal amount for the Great Lakes-St. Lawrence River Basin, excluding instream hydroelectric water use was 160,567 million liters per day (mld) or 44,493 million gallons per day (mgd). This total represents a five percent increase from the 2013 reported withdrawal amount total of 160,426 mld. Five and a half percent of the total reported amount withdrawn (9,282 mld) was consumed or otherwise lost to the basin.

Water withdrawals for all water use sectors, excluding the in-stream hydroelectric water use sector, are presented in the pie chart below. The 10 water use sectors are defined in Appendix C. Thermoelectric power production, industrial and public water supply are the primary water use sectors, (i.e., those withdrawing the largest volumes of water).

¹ An Olympic size swimming pool holds at least 2.5 million liters.



Water Withdrawals by Water Use Sector in mgd (excluding in-stream hydroelectric water use)

Figure 2.

The Lake Michigan watershed has the greatest withdrawal, followed by Lakes Ontario and Erie, respectively. The graph below shows withdrawals by watershed broken down by water source, e.g., Great Lakes surface water (GLSW), other surface water (OSW) and groundwater (GW).

Water Withdrawals by Watershed in mgd (excludes in-stream hydroelectric water use)





Figure 3.

The pie chart below shows total withdrawals portioned by jurisdiction, excluding in-stream hydroelectric water use. Ontario, with a land area spanning five watersheds (i.e., lake basins) was the largest withdrawer of Great Lakes water at 66,267 mld or 39 percent of the total withdrawal amount. In contrast, Pennsylvania, with a land area of 1,316 square kilometers within the Lake Erie watershed, withdrew just 136 mld.



Water Withdrawals by Jurisdiction in mgd (excludes in-stream hydroelectric water use)

Figure 4.

Hydroelectric Power Generation

In the past years of reporting annual water use, withdrawals for hydroelectric purposes (both in-stream and off-stream) have been the largest single sector of use, typically representing more than 95 percent of the region's total water withdrawals. In-stream hydroelectric power production continues to be a major water use for the Great Lakes-St. Lawrence River region (e.g., New York produced more hydroelectric power than any other state east of the Rocky Mountains in 2011²). However, beginning in 2012, under the 2009 water use data collection and reporting protocols, the reporting of in-stream hydroelectric power production data is optional. In-stream hydroelectric power water use is not considered a withdrawal because the water remains in the water body and is not associated with water consumption.

The regional water use database is designed to receive and report on both 1) off-stream hydroelectric power production water withdrawals; and 2) in-stream hydroelectric water use data submitted by the states and provinces. For 2014, the regional database has incomplete data for the in-stream hydroelectric water use sector. Québec and Indiana did not report hydroelectric water use, although both jurisdictions reported it in the past.

² U.S. Energy Information Administration. 2012. http://www.eia.gov/state/?sid=NY

Off-stream hydroelectric is considered a withdrawal since the water is removed to a retention area or a reservoir that serves as a pump-storage system. This storage substantially increases the surface area of the water body, and in so doing, increases the evaporation expected from that body of water, resulting in a consumptive use. After being used, the water is returned to the original water source. Both off-stream and in-stream totals are presented in a chart format in each of the watershed and jurisdiction summaries contained in this report. In 2014, a withdrawal total of 155 mld was reported for the off-stream hydroelectric power production sector. This withdrawal amount was reported by the states of New York and Minnesota.

Diversions and Consumptive Uses

Diversions and consumptive uses of water are key components of the regional water use database. The Compact and Agreement definitions for diversions and consumptive use are presented in Appendix B. Consumptive uses and diversions (less return flow) reflect water not being returned (i.e., water loss) to the source watershed. These water use data are considered particularly informative for assessing the cumulative hydrologic effects of water use in the region.

The total reported 2014 diversion from the Great Lakes-St. Lawrence River Basin was 4,792 mld or 1,266 mgd. The majority (88% or 4,243 mld) of this amount was associated with the Illinois diversion, which takes water from Lake Michigan and discharges it into the Mississippi River watershed. The reported amount associated with the Illinois diversion is relatively constant with the 2013 reported amount of 4,240 mld. Smaller diversions throughout the region make up the balance of the total, and some of the diverted water is returned to the source watershed as return flow. There are a number of diversions into the Basin, including the Long Lac and Ogoki diversions (two incoming diversions from the Hudson Bay watershed into northern Lake Superior) which contributed 12,802 mld (3,382 mgd) to the entire Basin in 2014. This is an increase of 1,393 mld or 12 percent from the 2013 reported amount of 11,409 mld. This increase is well within the range of flow variability observed from 1944-2013. The flow from these diversions has ranged from 6,219 to 19,612 mld³. When conditions in the Long Lac and Nipigon (downstream of Ogoki) watersheds are wet, the diversions are often reduced, and water that otherwise would have been diverted into Lake Superior is instead directed through natural outlets that flow toward Hudson Bay. Conversely, when conditions are dry in the downstream watersheds, the diversion flow may be higher. Overall, the net diversion, i.e., incoming diversions minus outgoing diversions, is a gain of 8,029 mld⁴ (2,121mgd), meaning that more water is diverted into the Basin than is diverted out of the Basin.

Consumptive use is defined as that portion of the water withdrawn or withheld from the Basin that is lost or otherwise not returned to the Basin due to evaporation, incorporation into products, or other processes. Consumptive use is most often calculated by applying a consumptive use coefficient to the reported withdrawal amount. The current database framework documents the consumptive use coefficient used for each water withdrawal record and the percentage of the consumptive use amount that was determined through actual measurement. The total reported consumptive use for the Basin was 9,426 mld (2,490 mgd) - an increase of 598 mld or nearly 7 percent from the 2013 total consumptive use amount of 8,831 mld. The public water supply at 2,725 mld and industrial at 2,786 mld were primary contributors to the total consumptive use amount. At 3,967 mld (1,047 mgd), the Lake Michigan watershed had the largest consumptive use total among the five lake watersheds and the St. Lawrence River watershed. Taking in consideration both consumptive use and diversions, the reported water loss to the Basin was 1,397 mld

³ Information on the flow variability of the Long Lac and Ogoki diversions was provided by Ontario Power Generation.

⁴ The Great Lakes Regional Water Use Database records all incoming diversions with a negative sign and all outgoing diversions with a positive sign. This sign convention is different from what is used in the interim cumulative impact assessment, Appendix D.

(369 mgd) for the year 2014, which is a 56 percent decrease from the 2013 net water loss of 3,206 mld. Tables 1-3 summarize water withdrawals, diversions and consumptive uses at the regional scale.

| Watershed | | Withdr | awals | Diver | rsions | Consumptive | |
|--------------------|-----------|---------|-------|-----------|------------|-------------|-------|
| Watershed | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Lake Superior | 2,270 | 125,586 | 80 | 127,936 | 0 | -12,772 | 175 |
| Lake Michigan | 37,548 | 3,844 | 2,239 | 43,631 | 2 | 4,611 | 3,964 |
| Lake Huron | 26,535 | 67,186 | 350 | 94,072 | 158 | 0 | 535 |
| Lake Erie | 197,230 | 5,273 | 981 | 203,484 | 15,827 | -49 | 1,667 |
| Lake Ontario | 190,666 | 374,449 | 245 | 565,361 | -15,974 | 155 | 1,412 |
| St. Lawrence River | 609,125 | 200,115 | 414 | 809,655 | 0 | 26 | 1,673 |
| Total | 1,063,375 | 776,454 | 4,309 | 1,844,138 | 13 | -8,029 | 9,426 |

Table 3. Basin 2014 Water Use Data Summary by Watershed

In millions of liters per day; Water Sources: Great Lakes surface water (GLSW), other surface water (OSW) and groundwater (GW)

Table 4. Basin 2014 Water Use Data Summary by Sector

| Sector | | Withd | rawals | | Diver | rsions | Consumptive |
|---|-----------|---------|--------|-----------|------------|------------|-------------|
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 16,119 | 2,840 | 1,778 | 20,736 | 13 | 3,626 | 2,725 |
| Self-Supply Commercial & Institutional | 223 | 212 | 52 | 488 | 0 | 8 | 87 |
| Self-Supply Irrigation | 11 | 497 | 1,024 | 1,532 | 0 | 0 | 1,347 |
| Self-Supply Livestock | 2 | 339 | 242 | 583 | 0 | 0 | 75 |
| Self-Supply Industrial | 12,082 | 4,349 | 1,142 | 17,573 | 0 | 114 | 2,786 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 101,452 | 10,464 | 8 | 111,924 | 0 | 0 | 1,917 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 2,878 | 88 | 23 | 2,989 | 0 | 49 | 337 |
| Off-Stream Hydroelectric Power Production | 0 | 156 | 0 | 156 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 929,589 | 753,982 | 0 | 1,683,571 | 0 | -12,802 | 0 |
| Other Self Supply | 1,019 | 3,526 | 40 | 4,586 | 0 | 977 | 151 |
| Total | 1,063,375 | 776,454 | 4,309 | 1,844,138 | 13 | -8,029 | 9,426 |

In millions of liters per day; Water Sources: Great Lakes surface water (GLSW), other surface water (OSW) and groundwater (GW)

Table 5. Basin 2014 Water Use Data Summary by Jurisdiction (includes in-stream hydro)

| Jurisdiction | | Withdr | awals | | Diver | rsions | Consumptive |
|--------------|-----------|---------|-------|-----------|------------|------------|-------------|
| Juliouretion | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Illinois | 6,099 | 0 | 0 | 6,099 | 0 | 4,252 | 2 |
| Indiana | 6,964 | 258 | 396 | 7,618 | 0 | 312 | 1,599 |
| Michigan | 30,146 | 5,417 | 1,930 | 37,494 | 2 | 0 | 1,952 |
| Minnesota | 1,238 | 9,646 | 24 | 10,908 | 0 | 31 | 121 |
| New York | 490,140 | 311,491 | 111 | 801,742 | 0 | 169 | 942 |
| Ohio | 5,249 | 1,546 | 303 | 7,098 | 0 | -76 | 509 |
| Ontario | 507,599 | 444,908 | 796 | 953,302 | 11 | -12,802 | 1,366 |
| Pennsylvania | 117 | 8 | 12 | 138 | 0 | 0 | 23 |
| Québec | 2,977 | 1,481 | 292 | 4,751 | 0 | 12 | 1,565 |
| Wisconsin | 12,847 | 1,699 | 443 | 14,989 | 0 | 75 | 1,348 |
| Total | 1,063,375 | 776,454 | 4,309 | 1,844,138 | 13 | -8,029 | 9,426 |

In millions of liters per day; Water Sources: Great Lakes surface water (GLSW), other surface water (OSW) and groundwater (GW)

Lake Watershed Summaries

Lake Superior




Figure 5. Lake Superior Watershed

Overview of Watershed Characteristics

Lake Superior is the largest of the Great Lakes and the world's third-largest freshwater lake by volume, holding about 12,100 cubic kilometers (2,900 cubic miles) of water. Lake Superior can hold all the water in the other Great Lakes, plus three more Lake Eries.⁵ Its surface area is roughly the size of South Carolina, or approximately 82,103 square kilometers (31,700 square miles).

Basic Stats of Lake Superior

Length: 350 mi / 563 km

Breadth: 160 mi / 257 km

Elevation: 600 ft / 183 m

Depth: 483 ft / 147 m average, 1,330 ft / 406 m maximum

Volume: 2,900 cubic mi / 12,100 cubic km

Lake Surface Area: 31,700 square mi / 82,100 square km

Watershed Drainage Area: 49,300 square mi / 127,700 square km

Outlet: St. Marys River to Lake Huron

Retention / Replacement Time¹: 191 years

Population in the Watershed: United States 444,000; Canada 229,000. Total: 673,000

 $^{{}^{5}\!}Retention$ time is the calculated quantity expressing the mean time water spends in the lake

Minnesota Sea Grant. 2012. http://www.seagrant.umn.edu/superior/facts

Water Withdrawals

Four jurisdictions share the Lake Superior watershed – Michigan, Minnesota, Ontario and Wisconsin – which collectively withdrew3,770 mld of water, excluding in-stream hydroelectric water use (124,166 mld). This amount is a 0.8 percent decrease from the 2013 total withdrawal amount of 3,801 mld. Thermoelectric power production, once-through and recirculated cooling (2,192 mld) and industrial (1,061 mld) were the major water use sectors.

Other surface waters within the Lake Superior watershed were primarily used to generate electricity with in-stream hydroelectric. Excluding in-stream hydroelectric water use, 60 percent (2,271 mld) of the total reported withdrawal amount from the watershed came directly from Lake Superior. The remaining withdrawals came directly from other surface waters (38% or 1,420 mld) and groundwater (2% or 79 mld).

Water Diversions and Consumptive Uses

Reported water loss in the Lake Superior watershed totaled 208 mld (5 percent of the total withdrawal amount). It was comprised mainly of the aggregated industrial water diversion in Minnesota of 31 mld and the total watershed consumptive use for all four jurisdictions of 175 mld. Industrial use (112 mld), use for thermoelectric power, recirculated cooling (26 mld) and the public water supply (28 mld) sectors were the largest contributors, respectively, to the total consumptive use for the watershed.

Water gain (12,802 mld) in the Lake Superior watershed came from the historic Long Lac and Ogoki diversion in Northern Ontario. On average, these diversions into the basin together were about two times by volume larger than the Illinois diversion out of the Basin.

| Sector | | Withd | awals | | Diver | sions | Consumptive | |
|---|-------|---------|-------|---------|------------|------------|-------------|--|
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use | |
| Public Water Supply | 179 | 11 | 60 | 250 | 0 | 0 | 28 | |
| Self-Supply Commercial & Institutional | 2 | 5 | 0 | 7 | 0 | 0 | 1 | |
| Self-Supply Irrigation | 0 | 1 | 4 | 4 | 0 | 0 | 4 | |
| Self-Supply Livestock | 2 | 95 | 10 | 107 | 0 | 0 | 2 | |
| Self-Supply Industrial | 579 | 476 | 6 | 1,061 | 0 | 31 | 112 | |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 1,417 | 775 | 0 | 2,192 | 0 | 0 | 26 | |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 90 | 0 | 0 | 90 | 0 | 0 | 1 | |
| Off-Stream Hydroelectric Power Production | 0 | 58 | 0 | 58 | 0 | 0 | 0 | |
| In-Stream Hydroelectric Water Use | 0 | 124,166 | 0 | 124,166 | 0 | -12,802 | 0 | |
| Other Self Supply | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 2,270 | 125,586 | 80 | 127,936 | 0 | -12,770 | 175 | |

Table 6. Lake Superior Watershed 2014 Water Use Data Summary

In millions of liters per day

Lake Michigan




Figure 6. Lake Michigan Watershed

Overview of Watershed Characteristics

Lake Michigan is the only one of the Great Lakes entirely within the United States. It is the second largest of the Great Lakes by volume, holding about 4,918 cubic kilometers (1,180 cubic miles) of water. Its surface area is roughly the size of West Virginia, approximately 57,753 square kilometers (22,300 square miles). More than 12 million people call the Lake Michigan watershed home; about a third of the entire population of the Great Lakes-St. Lawrence River Basin lives in the Lake Michigan watershed.

Water Withdrawals

Four jurisdictions share the Lake Michigan watershed – Illinois, Indiana, Michigan, and Wisconsin – and collectively used 43,631 mld. This amount is a 2.5 percent decrease from the 2013 total withdrawal amount of 44,781

Basic Stats of Lake Michigan

Length: 307 mi / 494 km Breadth: 118 mi / 190 km Elevation: 577.5 ft / 176 m Depth: 279 ft / 85 m average, 923 ft / 281 m maximum Volume: 1,180 cubic mi / 4,918 cubic km Lake Surface Area: 22,300 square mi / 57,753 square km Watershed Drainage Area: 45,600 square mi / 118,095 square km Outlet: Straits of Mackinac to Lake Huron

Retention / Replacement Time: 62 years

Population in the Watershed: 12,052,743

mld. The primary water uses were thermoelectric power, both once-through and recirculated cooling (27,879 mld), industrial use (7,460 mld) and public water supply (5,965 mld). Lake Michigan (86% of total withdrawals or 37,548 mld) was the primary source of water withdrawals in the watershed.

Water Diversions and Consumptive Uses

Reported water loss in the Lake Michigan watershed, totaling 8,574 mld, represents 17 percent of total withdrawals. Total water loss was comprised mainly of the Illinois diversion (4,243 mld for public water supply and other purposes), additional small diversion totaling 367 mld and the total consumptive use of the four jurisdictions (3,964 mld). Industrial use (1,433 mld), irrigation (970 mld) and public water supply (300 mld) were the water use sectors that contribute the majority of the consumptive uses in the watershed.

| Sector | | Withdr | awals | | Diver | sions | Consumptive |
|---|--------|--------|-------|--------|------------|------------|-------------|
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 5,014 | 81 | 870 | 5,965 | 2 | 3,532 | 300 |
| Self-Supply Commercial & Institutional | 14 | 18 | 34 | 66 | 0 | 9 | 7 |
| Self-Supply Irrigation | 1 | 188 | 921 | 1,111 | 0 | 0 | 970 |
| Self-Supply Livestock | 0 | 65 | 101 | 165 | 0 | 0 | 31 |
| Self-Supply Industrial | 6,051 | 1,144 | 265 | 7,460 | 0 | 82 | 1,433 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 24,971 | 2,259 | 6 | 27,236 | 0 | 0 | 1,085 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 539 | 84 | 19 | 643 | 0 | 49 | 137 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Self Supply | 958 | 4 | 23 | 985 | 0 | 940 | 1 |
| Total | 37,548 | 3,844 | 2,239 | 43,631 | 2 | 4,611 | 3,964 |

Table 7. Lake Michigan Watershed 2014 Water Use Data Summary

In millions of liters per day

Lake Huron




Figure 7. Lake Huron Watershed

Overview of Watershed Characteristics

By surface area, Lake Huron is the second-largest of the Great Lakes. It covers 59,600 square kilometers (23,000 square miles), making it the third-largest fresh water lake on Earth. By volume however, Lake Huron is only the third largest of the Great Lakes.

Water Withdrawals

Two jurisdictions – Michigan and Ontario – share the watershed and collectively used 32,503 mld of the water, excluding in-stream hydroelectric water use (61,569 mld). This is a 6 percent increase from the 2013 water withdrawal amount of 830,571mld.

The primary water uses were industrial use (1,428 mld), thermoelectric power, once-through cooling (29,923 mld) and public water supply (881 mld). Excluding in-stream

Basic Stats of Lake Huron

Length: 206 mi / 332 km

Breadth: 183 mi / 295 km

Elevation: 577.5 ft / 176 m

Depth: 195 ft / 59 m average, 750 ft / 229 m maximum

Volume: 849 cubic mi / 3,538 cubic km Lake Surface Area: 23,000 square mi / 59,565 square km

Watershed Drainage Area: 50,700 square mi / 131,303 square km

Outlet: St. Clair River to Lake Erie

Retention / Replacement Time: 21 years

Population in the Watershed: United States 1,483,872; Canada 1,476,487. Total: 2,960,359 hydroelectric water use, Lake Huron surface water was the source of 82 percent of the total withdrawals in the watershed.

Water Diversions and Consumptive Uses

Reported water loss to the Lake Huron watershed was 693 mld, which represented nearly two percent of the total withdrawal amount. This total includes an intrabasin transfer of 158 mld for public water supply in Ontario. While hydrologically this intrabasin transfer remained in the Great Lakes-St. Lawrence River Basin, it represented a loss to the Lake Huron watershed and a net gain to the Lake Erie watershed. Public water supply (108 mld) and thermoelectric power production (250 mld) made up the majority (67%) of the consumptive uses in the watershed.

| Sector | | Withdr | awals | | Diver | sions | Consumptive |
|---|--------|--------|-------|--------|------------|------------|-------------|
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 561 | 162 | 158 | 881 | 158 | 0 | 108 |
| Self-Supply Commercial & Institutional | 5 | 12 | 1 | 18 | 0 | 0 | 2 |
| Self-Supply Irrigation | 4 | 56 | 40 | 100 | 0 | 0 | 90 |
| Self-Supply Livestock | 0 | 44 | 82 | 125 | 0 | 0 | 18 |
| Self-Supply Industrial | 914 | 449 | 65 | 1,428 | 0 | 0 | 67 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 25,050 | 4,874 | 0 | 29,923 | 0 | 0 | 248 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 1 | 3 | 4 | 0 | 0 | 2 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 61,569 | 0 | 61,569 | 0 | 0 | 0 |
| Other Self Supply | 2 | 21 | 0 | 23 | 0 | 0 | 0 |
| Total | 26,535 | 67,186 | 350 | 94,072 | 158 | 0 | 535 |

Table 8. Lake Huron Watershed 2014 Water Use Data Summary

In millions of liters per day

Lake Erie




Figure 8. Lake Erie Watershed

Overview of Watershed Characteristics

By surface area, Lake Erie is the 12th largest freshwater lake in the world. The shallowest of the Great Lakes, it has an average depth of 19 meters and a maximum depth of 64 meters. The lake holds about 4,863 cubic kilometers (116 cubic miles) of water. Lake Erie is warmer than the other Great Lakes, which also helps make it the most productive. However, its size also makes it more ecologically sensitive than the other Great Lakes.

The watershed is home to more than 12.5 million people, representing more than one-third of the entire population of the Great Lakes-St. Lawrence River Basin.

Water Withdrawals

Six jurisdictions – Indiana, Michigan, New York, Ohio, Ontario and Pennsylvania – share the watershed and

Basic Stats of Lake Erie

Length: 241 mi / 388 km

Breadth: 57 mi / 92 km

Elevation: 569.2 ft / 173.5 m

Depth: 62 ft / 19 m average, 210 ft / 64 m maximum

Volume: 116 cubic mi / 483 cubic km

Lake Surface Area: 9,910 square mi / 25,655 square km

Watershed Drainage Area: 22,700 square mi / 58,788 square km

Outlets: Niagara River and Welland Canal

Retention/Replacement Time: 2.7 years

Population in the Watershed: United States, est. 10,640,671; Canada est. 1,892,306. Total: est. 12,532,977

collectively used 34,063 mld of the water, excluding in-stream hydroelectric water use, which accounted for 169,420 mld. This amount is a four percent decrease from the 2013 total withdrawal amount of 35,318 mld. Aside from water used for hydroelectric power generation purposes, the primary water uses were thermoelectric power, both once-through and recirculated cooling (23,586 mld), public water supply (6,004 mld) and industrial use (4,149 mld).

Lake Erie surface water was the source of 88 percent of the total withdrawals in the watershed. However, other surface water and groundwater were the only sources of Indiana's water use, primarily supporting the public water supply sector.

Water Diversions and Consumptive Uses

Reported water loss in the Lake Erie watershed totaled 17,485 mld. This amount includes an aggregate diversion (going into the Lake Erie watershed) of 26 mld, an aggregate intrabasin diversion of 15,827 mld, and a total consumptive use of 1,667 mld. The largest intrabasin diversion is the Welland Canal for other self supply, navigation purposes (15,977 mld). The Welland Canal was constructed in 1830 as a ship canal in Ontario, Canada, connecting Lake Erie to Lake Ontario. The major consumptive uses were for public water supply (774 mld) and industrial uses (344 mld).

| | | | | 5 | | | |
|---|---------|--------|-------|---------|------------|------------|-------------|
| Sector | | Withdr | awals | | Diver | sions | Consumptive |
| 5000 | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 4,928 | 674 | 402 | 6,004 | -150 | 77 | 774 |
| Self-Supply Commercial & Institutional | 1 | 10 | 8 | 20 | 0 | 0 | 2 |
| Self-Supply Irrigation | 5 | 136 | 53 | 194 | 0 | 0 | 174 |
| Self-Supply Livestock | 0 | 27 | 23 | 50 | 0 | -1 | 13 |
| Self-Supply Industrial | 2,735 | 935 | 480 | 4,149 | 0 | 0 | 344 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 21,682 | 1,276 | 1 | 22,960 | 0 | 0 | 236 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 624 | 2 | 0 | 626 | 0 | 0 | 116 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 167,239 | 2,181 | 0 | 169,420 | 0 | 0 | 0 |
| Other Self Supply | 14 | 32 | 14 | 61 | 15,977 | -84 | 8 |
| Total | 197,230 | 5,273 | 981 | 203,484 | 15,827 | -7 | 1,667 |

Table 9. Lake Erie Watershed 2014 Water Use Data Summary

In millions of liters per day

Lake Ontario




Figure 97. Lake Ontario Watershed

Overview of Watershed Characteristics

Lake Ontario is the easternmost of the Great Lakes and the smallest in surface area covering 18,960 square kilometers (7,340 square miles). It is extremely deep (e.g., 244 m at maximum) in some areas and exceeds Lake Erie in volume (1,639 cubic kilometers, 393 cubic miles). By surface area, it is the 14th largest lake in the world.

Water Withdrawals

Three jurisdictions – New York, Ontario and Pennsylvania – share the watershed and collectively used 40,101 mld of the water, excluding in-stream hydroelectric water use, which accounted for 525,260 mld. This amount is a 0.7 percent increase from the 2013 withdrawal amount of 39,819 mld. Aside from withdrawals for hydroelectric power generation purposes, the primary water uses were for public water supply (3,684 mld), other self-supply uses (2,943 mld) and

Basic Stats of Lake Ontario

Length: 193 mi / 311 km

Breadth: 53 mi / 85 km

Elevation: 243.3 ft / 74.2 m

Depth: 283 ft / 86 m average, 802 ft / 244 m maximum

Volume: 393 cubic mi / 1,639 cubic km

Lake Surface Area: 7,340 square mi / 19,009 square km

Watershed Drainage Area: 23,400 square mi / 60,601 square km

Outlet: St. Lawrence River to the Atlantic Ocean

Retention / Replacement Time: 6 years

Population in the Watershed: United States, est. 2,856,360; Canada est. 2,835,818. Total: est.5,692,178 thermoelectric power generation, both once-through and recirculated cooling (31,032 mld).

Lake Ontario surface water was the source for 85 percent of the total withdrawals in the watershed. It was the source for most of the water use sectors, except for irrigation and livestock where other surface water in the watershed was the predominant source.

Water Diversions and Consumptive Uses

Reported water loss in the Lake Ontario watershed totaled 1,563 mld. This amount includes diversions totaling 156 mld, an intrabasin diversion of 3 mld and a combined consumptive use amount of 1,415 mld. The major consumptive uses were from public water supply (455 mld), thermoelectric power production (401 mld) and industrial uses (268 mld).

Water gained (15,974 mld) in the Lake Ontario watershed came from the Welland Canal, which diverts water from the Lake Erie watershed for navigation purposes.

| Sector | | Withd | awals | | Diver | sions | Consumptive |
|---|---------|---------|-------|---------|------------|------------|-------------|
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 2,888 | 733 | 63 | 3,684 | 3 | 35 | 455 |
| Self-Supply Commercial & Institutional | 200 | 143 | 4 | 346 | 0 | 0 | 65 |
| Self-Supply Irrigation | 0 | 97 | 4 | 101 | 0 | 0 | 91 |
| Self-Supply Livestock | 0 | 78 | 9 | 87 | 0 | 0 | 8 |
| Self-Supply Industrial | 1,119 | 526 | 164 | 1,809 | 0 | 0 | 268 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 28,131 | 1,273 | 1 | 29,405 | 0 | 0 | 319 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 1,625 | 1 | 0 | 1,626 | 0 | 0 | 82 |
| Off-Stream Hydroelectric Power Production | 0 | 98 | 0 | 98 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 156,703 | 368,557 | 0 | 525,260 | 0 | 0 | 0 |
| Other Self Supply | 2 | 2,941 | 0 | 2,943 | -15,977 | 121 | 123 |
| Total | 190,666 | 374,449 | 245 | 565,361 | -15,974 | 156 | 1,412 |

Table 10. Lake Ontario Watershed 2014 Water Use Data Summary

In millions of liters per day

St. Lawrence River




Figure 10. St. Lawrence River Watershed

Overview of Watershed Characteristics

Running 1,198 kilometers (744 miles) in length, the St. Lawrence River is considered a major river of North America. Mostly located in the province of Québec, it links the Great Lakes to the Atlantic Ocean.

Water Withdrawals

Three jurisdictions – New York, Ontario and Québec – share the watershed and collectively used 6,498 mld of the water, excluding in-stream hydroelectric water use, which accounted for 803,157 mld. This amount is a 6 percent increase from the 2013 withdrawal total of 6,136 mld. Aside from hydroelectric, the primary water uses were public water supply (3,951 mld), and industrial use (1,666 mld).

Basic Stats of the St. Lawrence River

Length: 744 mi / 1,197 km Elevation: 245 ft/74.7 m at the source and 0 ft/0 m at the mouth

Average Annual Flow (Montréal): 7,660 cubic meters/second

Volume: 393 cubic mi / 1,639 cubic km

Watershed Drainage Area: 519,000 square mi / 1,344,200 square km Outlet: Gulf of St. Lawrence/ Atlantic Ocean

St. Lawrence River surface water was the source for over half (54%) of the watershed's total withdrawal amount. Other surface water within the St. Lawrence River watershed accounted for 40 percent of the total. The remaining portion of the total withdrawal amount (6%) came from groundwater sources.

Water Diversions and Consumptive Uses

Water loss in the St. Lawrence River watershed totaled 1,700 mld. This total includes a diversion amount of 25 mld for public supply purposes in New York and Québec and a combined consumptive use amount of 1,673 mld. The largest consumptive uses were the public water supply sector at 1,060 mld and industrial at 562 mld.

| Sector | | Withdr | awals | | Diver | sions | Consumptive | | |
|---|---------|---------|-------|---------|------------|------------|-------------|--|--|
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use | | |
| Public Water Supply | 2,549 | 1,178 | 225 | 3,951 | 0 | 25 | 1,060 | | |
| Self-Supply Commercial & Institutional | 2 | 25 | 5 | 31 | 0 | 0 | 9 | | |
| Self-Supply Irrigation | 1 | 19 | 2 | 22 | 0 | 0 | 19 | | |
| Self-Supply Livestock | 0 | 30 | 18 | 48 | 0 | 0 | 2 | | |
| Self-Supply Industrial | 684 | 819 | 162 | 1,666 | 0 | 0 | 562 | | |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 200 | 7 | 0 | 207 | 0 | 0 | 2 | | |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| In-Stream Hydroelectric Water Use | 605,647 | 197,510 | 0 | 803,157 | 0 | 0 | 0 | | |
| Other Self Supply | 43 | 528 | 3 | 574 | 0 | 0 | 19 | | |
| Total | 609,125 | 200,115 | 414 | 809,655 | 0 | 25 | 1,673 | | |

Table 11. St. Lawrence River Watershed 2014 Water Use Data Summary

In millions of gallons per day

Jurisdiction Reports

Illinois

The Illinois portion of the Lake Michigan watershed is only about 259 square kilometers, which accounts for less than 0.2 percent of the total area of the state. The Lake Michigan coastline of Illinois is 101 kilometers long, which is a small fraction of the 2,639 kilometers that make up the total Lake Michigan shoreline. Despite its relatively small size, the Illinois Lake Michigan service area is home to half of the total population of Illinois and the lake itself is the largest public drinking water supply in the state, serving nearly 7 million people.

The total withdrawal amount from the Basin for Illinois in 2014 was 6,099 mld, an 8 percent decrease from 2013 (6,693 mld). The largest uses of reported water were public water supply at 3,304 mld (nearly half of the total withdrawal amount) and thermoelectric power production, once-through cooling at 1,741 mld (29% of the total withdrawal amount). The primary source for all withdrawals was Lake Michigan.

The Illinois Diversion, which takes water from Lake Michigan and discharges it into the Mississippi River watershed, is comprised of three elements: public water supply; stormwater runoff; and support for control structures for navigation and discretionary diversion for other purposes such as low flow augmentation and water quality enhancement. The amount of water diverted for public water supply and self-supply commercial and institutional was 3,312 mld. The diversion amount supporting other uses (i.e., discretionary diversion) was 940 mld.

Data collected for this report came from multiple sources: Illinois Department of Natural Resources, Illinois State Water Survey and the Metropolitan Water Reclamation District of Greater Chicagoland. These data were generated with a 100 percent reporting compliance from permitted water withdrawal facilities.

| Sector | | Withdr | awals | | Diver | sions | Consumptive |
|---|-------|--------|-------|-------|------------|------------|-------------|
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 3,304 | 0 | 0 | 3,304 | 0 | 3,304 | 0 |
| Self-Supply Commercial & Institutional | 8 | 0 | 0 | 8 | 0 | 8 | 0 |
| Self-Supply Irrigation | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Self-Supply Livestock | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Self-Supply Industrial | 105 | 0 | 0 | 105 | 0 | 0 | 1 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 1,741 | 0 | 0 | 1,741 | 0 | 0 | 0 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Self Supply | 940 | 0 | 0 | 940 | 0 | 940 | 0 |
| Total | 6,099 | 0 | 0 | 6,099 | 0 | 4,252 | 2 |

Table 12. Illinois 2014 Water Use Data Summary

In millions of liters per day

Indiana

The state of Indiana relies on the water resources of the Lake Michigan and Lake Erie watersheds. Indiana's portion of Lake Michigan encompasses a total of 624 square kilometers. Four Indiana counties lie partially within the Lake Michigan watershed, but three of these four counties (Lake, Porter and LaPorte) constitute more than 99.5 percent of the land area in the watershed. Abundant freshwater from Lake Michigan has promoted the development of an extensive urban and industrial belt along the Indiana coast of Lake Michigan. Water supplies in Indiana's non-coastal counties in the Lake Michigan watershed are drawn primarily from groundwater sources.

Indiana shares a portion of the Maumee River watershed that flows into Lake Erie. The Maumee River watershed encompasses a total of 3,323 square kilometers of northeast Indiana. Six Indiana counties lie partially within this watershed. The largest withdrawals come from the surface waters of the St. Joseph (a major tributary within the Maumee watershed) and the Maumee River, used for public supply and industrial purposes. Groundwater withdrawals in the Maumee River watershed are used primarily for public and domestic water supply and dewatering for industrial purposes.⁶

In 2014, the total reported water withdrawal amount from the Basin for Indiana was 7,618 mld. The largest uses were industrial (5,838 mld), thermoelectric power (940 mld) and public water supply (641 mld).

The total reported diversion amount for Indiana was 312 mld. Because a small, 168 square kilometer portion of Indiana drains into the Illinois River (as a result of the Illinois Diversion), water transferred from the Lake Michigan watershed into this area is considered a diversion of water from the Great Lakes-St. Lawrence River Basin. The majority of reported diversions for Indiana (201 mld) were distributed for public supply purposes from Lake Michigan surface water and discharged to the "Illinois Diversion" area, with about 4 mld reported as a diversion from groundwater for public supply. The industrial sector was responsible for about 83 mld of the reported diversion from the Lake Michigan watershed to the Illinois River.

For the Lake Erie watershed, a portion of the town of Fort Wayne's public water supply distribution system is located in the Upper Wabash watershed. The amount of water (about 26 mld primarily from other surface water with a small portion from groundwater) distributed through that portion of the system was reported as a diversion from the Lake Erie watershed. Consumptive use totaled 1,599 mld, with the industrial sector in the Lake Michigan watershed (1,299 mld or 81%) as the primary contributor to the total.

Data collected for this report came from the Indiana Department of Natural Resources. These data were generated with reporting compliance rates from permitted water withdrawal facilities ranging from 75 to 100 percent depending on the water use sector. Withdrawals and consumptive uses are not estimated for facilities that did not report.

⁶ Indiana Dept. of Natural Resources. 1996. http://www.in.gov/dnr/water/files/lakemich_basinsums.pdf http://www.in.gov/dnr/water/files/maumee_basinsums.pdf

| | | | 5 | | | | |
|---|-------|--------|-------|-------|------------|------------|-------------|
| Sector | | Withdr | awals | | Diver | sions | Consumptive |
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 343 | 125 | 173 | 641 | 0 | 230 | 78 |
| Self-Supply Commercial & Institutional | 0 | 0 | 5 | 5 | 0 | 0 | 1 |
| Self-Supply Irrigation | 0 | 24 | 152 | 175 | 0 | 0 | 157 |
| Self-Supply Livestock | 0 | 4 | 10 | 14 | 0 | 0 | 7 |
| Self-Supply Industrial | 5,773 | 25 | 40 | 5,838 | 0 | 82 | 1,299 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 796 | 0 | 1 | 797 | 0 | 0 | 16 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 52 | 79 | 13 | 144 | 0 | 0 | 42 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Self Supply | 0 | 0 | 3 | 3 | 0 | 0 | 0 |
| Total | 6,964 | 258 | 396 | 7,618 | 0 | 312 | 1,599 |

Table 13. Indiana 2014 Water Use Data Summary

In millions of liters per day

Michigan

Home to more than 9.8 million people, Michigan borders four of the Great Lakes (Superior, Michigan, Huron and Erie). Some unique features of Michigan include:

- 147,686 square kilometers of land area in two peninsulas (105,109 square kilometers in the Lower Peninsula and 42,576 square kilometers in the Upper Peninsula);
- Virtually the entire land area of the state lies within the Great Lakes basin;
- 99,909 square kilometers of Great Lakes water area; and
- 5,031 kilometers of Great Lakes shoreline (more fresh water coastline than any other state).⁷

In 2014, the total reported water withdrawal amount from the Basin for Michigan was 37,494 mld, a decrease of 2 percent from the 2013 total water withdrawal amount of 38,271 mld. The largest use was thermoelectric power production, once-through and recirculated cooling, totaling 29,942 mld or nearly 80 percent of the total withdrawal amount. The Great Lakes proper were the largest source for withdrawals at 80 percent of the total. Nearly half of the total water withdrawal amount (18,223 mld, 49%) came from the Lake Erie watershed, mainly used for thermoelectric power production. Forty one percent of total withdrawal amount (15,248 mld) came from the Lake Michigan watershed, followed by the Lake Huron watershed at 3,131 mld (8%) and the Lake Superior watershed at 890 mld (2%).

Michigan reported no diversions. The total amount of consumptive use was 1,952 mld (5% of the total withdrawal amount), with self-supply irrigation being the largest contributor to consumptive use at 869 mld.

Data collected for this report came from multiple sources: Michigan Department of Environmental Quality and the Michigan Department of Agriculture and Rural Development. These data were generated with estimated reporting compliance rates ranging from 85 to 99 percent of total water use reporters, depending on the water use sector. Withdrawals and consumptive uses are not estimated for facilities not in compliance with reporting for most water use sectors except for self-supply livestock which was partially estimated by the state agency. This sector had 40 percent reporting compliance rate due to a lag in reporting within the agricultural community and the omission of aquaculture facilities from agency reporting.

⁷ Michigan Dept. of Transportation. http://www.michigan.gov/mdot/0,4616,7-151-9622_11033_11151-67959--,00.html

| 0 | | | | | | | |
|---|--------|--------|-------|--------|------------|------------|-------------|
| Sector | | Withdr | awals | | Diver | sions | Consumptive |
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 2,938 | 69 | 752 | 3,759 | 2 | 0 | 470 |
| Self-Supply Commercial & Institutional | 0 | 10 | 24 | 34 | 0 | 0 | 4 |
| Self-Supply Irrigation | 1 | 237 | 727 | 965 | 0 | 0 | 869 |
| Self-Supply Livestock | 0 | 84 | 78 | 162 | 0 | 0 | 26 |
| Self-Supply Industrial | 1,213 | 1,069 | 314 | 2,597 | 0 | 0 | 260 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 25,370 | 3,926 | 6 | 29,303 | 0 | 0 | 184 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 622 | 7 | 10 | 640 | 0 | 0 | 140 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Self Supply | 2 | 14 | 19 | 34 | 0 | 0 | 0 |
| Total | 30,146 | 5,417 | 1,930 | 37,494 | 2 | 0 | 1,952 |

Table 14. Michigan 2014 Water Use Data Summary

In millions of liters per day

Minnesota

The Minnesota part of the Lake Superior watershed encompasses approximately 16,058 square kilometers. Major river watersheds in the basin include the Cloquet, Nemadji and St. Louis River systems, as well as the north shore tributaries to Lake Superior.⁸

Excluding in-stream hydroelectric water use (8,747 mld), the total withdrawal amount from the Basin for Minnesota was 2,161 mld, a decrease of 2 percent from the total withdrawal amount for 2013 (2,203 mld). The major water use sectors include industrial at 822 mld and thermoelectric power production, once-through cooling at 1,137 mld. These water use sectors were about equally supplied by Lake Superior and other surface water within the Lake Superior watershed.

The total reported diversion amount of 31 mld was for industrial purposes. Total consumptive use was 121 mld, which is relatively constant compared to the 2013 total consumptive use of 125 mld. The majority of that amount (82 mld) was for industrial purposes.

The water use data was provided by the Minnesota Department of Natural Resources which collected measured water use data from water withdrawal permit holders with a 100 percent reporting compliance from permitted water withdrawal facilities.

| 5 | | | | | | | | | |
|---|-------|-------|--------|--------|------------|------------|-------------|--|--|
| Sector | | Withd | rawals | | Diver | rsions | Consumptive | | |
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use | | |
| Public Water Supply | 109 | 7 | 22 | 138 | 0 | 0 | 14 | | |
| Self-Supply Commercial & Institutional | 2 | 2 | 0 | 4 | 0 | 0 | 0 | | |
| Self-Supply Irrigation | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| Self-Supply Livestock | 2 | 0 | 0 | 2 | 0 | 0 | 2 | | |
| Self-Supply Industrial | 474 | 346 | 2 | 822 | 0 | 31 | 82 | | |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 650 | 486 | 0 | 1,137 | 0 | 0 | 23 | | |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| Off-Stream Hydroelectric Power Production | 0 | 58 | 0 | 58 | 0 | 0 | 0 | | |
| In-Stream Hydroelectric Water Use | 0 | 8,747 | 0 | 8,747 | 0 | 0 | 0 | | |
| Other Self Supply | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| Total | 1,238 | 9,646 | 24 | 10,908 | 0 | 31 | 121 | | |

Table 15. Minnesota 2014 Water Use Data Summary

In millions of liters per day

⁸ Minnesota Pollution Control Agency. 2013. http://www.pca.state.mn.us/index.php/water/water-types-and-programs/surface-water/basins/lake-superior-basin/index.html

New York

Approximately 80 percent of New York State's fresh surface water, over 1,127 kilometers of shoreline and nearly 48 percent of New York lands are contained in the drainage watersheds of Lake Erie, Lake Ontario and the St. Lawrence River, which includes the Lake Champlain/Lake George watersheds. More than four million New Yorkers depend on the fresh water of these watersheds for drinking water, and hundreds of miles of waterways and border waters for navigation.⁹

Excluding in-stream hydroelectric water use (787,790 mld), the total withdrawal amount from the Basin for New York was 13,952 mld, a 4 percent decrease from 2013 (14,562 mld). The Lake Ontario watershed was the source of the majority of New York's water withdrawals at 11,060 mld or 79 percent of the total withdrawal amount. Thermoelectric power production (both once-through and recirculated cooling) at 7,731 mld represented nearly 55 percent of the total withdrawal amount; public water supply (1,796 mld) represented nearly 13 percent of the total; and industrial (1,122 mld) represented 8 percent of the total. For the Lake Erie and Lake Ontario watersheds, Great Lakes surface water was the primary source of water, when in-stream hydroelectric is excluded. For the St. Lawrence River watershed, other surface water was the primary source of water, when in-stream hydroelectric is excluded.

The 2014 total diversion amount for New York was 169 mld of which 48 mld was for public supply and 121 mld for other self-supply purposes. The total consumptive use amount was 942 mld. The largest consumptive use was attributed to industrial purposes at 241 mld.

The water use data was provided by the New York State Department of Environmental Conservation. The data collected was metered and estimated water use. Reporting compliance varies among the water use sectors from 56 percent for the irrigation sector to 100 percent for the hydroelectric power sector. New York State does not estimate the water use for facilities that did not report their use.

| 5 | | | | | | | | | |
|---|---------|---------|--------|---------|------------|------------|-------------|--|--|
| Sector | | Withd | rawals | | Diver | sions | Consumptive | | |
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use | | |
| Public Water Supply | 1,112 | 613 | 71 | 1,796 | 0 | 48 | 22 | | |
| Self-Supply Commercial & Institutional | 0 | 159 | 6 | 166 | 0 | 0 | 3 | | |
| Self-Supply Irrigation | 0 | 109 | 5 | 114 | 0 | 0 | 10 | | |
| Self-Supply Livestock | 0 | 87 | 1 | 88 | 0 | 0 | 1 | | |
| Self-Supply Industrial | 604 | 495 | 23 | 1,122 | 0 | 0 | 24 | | |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 5,280 | 824 | 1 | 6,105 | 0 | 0 | 12 | | |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 1,625 | 1 | 0 | 1,626 | 0 | 0 | 8 | | |
| Off-Stream Hydroelectric Power Production | 0 | 98 | 0 | 98 | 0 | 0 | | | |
| In-Stream Hydroelectric Water Use | 481,520 | 306,270 | 0 | 787,790 | 0 | 0 | | | |
| Other Self Supply | 0 | 2,833 | 4 | 2,838 | 0 | 121 | 12 | | |
| Total | 490,140 | 311,491 | 111 | 801,742 | 0 | 169 | 94 | | |

Table 16. New York 2014 Water Use Data Summary

In millions of liters per day

⁹ Great Lakes Basin Advisory Council. 2013. Our Great Lakes Water Resources: Conserving and Protecting Our Water Today for Use Tomorrow Final Report. http://www.dec.ny.gov/docs/regions_pdf/glbacfrpt.pdf

Ohio

Ohio's portion of the Lake Erie watershed drains 30,171 square kilometers and is home to 4.65 million people. Toledo, Sandusky and Cleveland are some of the communities that dot Ohio's 502 kilometer-long shoreline. Agricultural row crops account for 59 percent of the land use in the Ohio watersheds draining to Lake Erie, followed by urban residential and commercial land use at a combined 16 percent. Another 16 percent were from forested lands and wetlands, combined with pasture land making up 5 percent of total land use.¹⁰

The 2014 total reported withdrawal amount from the Basin for Ohio was 7,098 mld, a 3 percent decrease from the total withdrawal amount for 2013 (7,310 mld). Primary water use sectors included thermoelectric power production (once-through and recirculated cooling) at 4,164 mld, representing 59 percent of total withdrawal amount; public water supply (1,996 mld), representing 28 percent; and industrial (797 mld), representing 11 percent. The source for 74 percent of the total withdrawal amount was Lake Erie. However, within the irrigation and industrial water use sectors, other surface water was the primary source at 90 percent and 52 percent of the total withdrawal amount, respectively.

Overall, 49 mld was diverted out of the Lake Erie watershed, all for public water supply purposes, of which 42 mld was returned to the watershed after use, resulting in a net diversion of 8 mld. This was offset by incoming diversions totaling 83 mld, primarily for other self-supply purposes, resulting in a net diversion of 75 mld into the Lake Erie watershed. Total consumptive use was 509 mld. Fifty nine percent of the total consumptive use was attributed to the public water supply sector.

The water use data was provided by the Ohio Department of Natural Resources, Division of Water Resources with a 100 percent reporting compliance from every water use sector.

| Sector | | Withdr | awals | | Diver | sions | Consumptive Use |
|---|-------|--------|-------|-------|------------|------------|--------------------|
| | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | |
| Public Water Supply | 1,520 | 375 | 101 | 1,996 | 0 | 8 | 299 |
| Self-Supply Commercial & Institutional | 1 | 0 | 0 | 1 | 0 | 0 | 0 |
| Self-Supply Irrigation | 2 | 102 | 9 | 113 | 0 | 0 | 101 |
| Self-Supply Livestock | 0 | 0 | 2 | 2 | 0 | 0 | 2 |
| Self-Supply Industrial | 194 | 415 | 188 | 797 | 0 | 0 | 17 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 3,086 | 637 | 0 | 3,723 | 0 | 0 | 37 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 441 | 0 | 0 | 441 | 0 | 0 | 44 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use* | 0 | 0 | 0 | 0 | | | 0 |
| Other Self Supply | 5 | 17 | 3 | 25 | 0 | -83 | 8 |
| Total | 5,249 | 1,546 | 303 | 7,098 | 0 | -75 | 509 |

Table 17. Ohio 2014 Water Use Data Summary

In millions of liters per day

Water Sources: Great Lakes surface water (GLSW), other surface water (OSW) and groundwater (GW) **No data was provided for this sector.*

¹⁰ Ohio Environmental Protection Agency. 2010. Ohio Lake Erie Phosphorus Task Force Final Report.

http://www.epa.ohio.gov/portals/35/lakeerie/ptaskforce/Task_Force_Final_Report_April_2010.pdf

Ontario

More than 98 percent of Ontario residents (more than 12 million people) live within the Great Lakes-St. Lawrence River Basin. Most live along the coast in eight of Canada's 20 largest cities, which include Toronto, Hamilton, Windsor and Sarnia.¹¹ Ontario's portion of the Great Lakes forms the longest freshwater coastline in the world stretching more than 6,800 miles (11,000 kilometers)¹² across five major watersheds in the Great Lakes-St. Lawrence River system: Lake Superior, Lake Huron, Lake Erie, Lake Ontario and the St. Lawrence River watersheds.

For this report, 2013 data was used for most all water use sectors except for the diversion amounts related to the public water supply sector for which 2009 data were used. Excluding in-stream hydroelectric water use (reported amount of 887,034 mld), the total water withdrawal amount from the Basin was approximately 66,267 mld. The three largest water use categories were thermoelectric power (once-through cooling) at 56,422 mld or 85 percent of the total withdrawal amount; public supply at 4,370 mld; and industrial at 4,382 mld. Except for the Lake Superior and St. Lawrence River watersheds, where other surface water was the primary source for withdrawals, the primary source for withdrawals came from Great Lakes surface water.

No diversions from the Great Lakes-St. Lawrence River Basin were reported for Ontario, while diversions into the basin were approximately 12,802 mld. The total consumptive use amount was approximately 1,366 mld. Three water use sectors, representing the largest consumptive uses, included thermoelectric power at 508 mld, public water supply at 525 mld and industrial at 289 mld. Ontario reported intrabasin diversions totaling 16,138 mld as summarized in Table 17.

These data were collected primarily through the provincial water taking and reporting system. Additional estimates were provided by water use sector to capture water use that was not reported. Reporting data varied among water use sectors from 88 percent for the industrial sector to 100 percent for thermoelectric power production (once-through cooling).

¹¹ Ontario Ministry of Natural Resources. 2012. http://www.mnr.gov.on.ca/en/Business/GreatLakes/2ColumnSubPage/STEL02_173888.html ¹² Ontario Ministry of the Environment. 2012,

http://www.ene.gov.on.ca/stdprodconsume/groups/lr/@ene/@resources/documents/resource/stdprod_096933.pdf

| | | | 5 | | | | |
|---|---------|---------|-------|---------|------------|------------|-------------|
| Sector | | Withdra | awals | | Diver | sions | Consumptive |
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 3,380 | 725 | 265 | 4,370 | 11 | 0 | 525 |
| Self-Supply Commercial & Institutional | 205 | 19 | 6 | 230 | 0 | 0 | 34 |
| Self-Supply Irrigation | 6 | 5 | 0 | 10 | 0 | 0 | 9 |
| Self-Supply Livestock | 0 | 99 | 90 | 189 | 0 | 0 | 2 |
| Self-Supply Industrial | 3,171 | 783 | 428 | 4,382 | 0 | 0 | 289 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 52,757 | 3,665 | 0 | 56,422 | 0 | 0 | 508 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 448,069 | 438,965 | 0 | 887,034 | 0 | -12,802 | 0 |
| Other Self Supply | 11 | 646 | 7 | 664 | 0 | 0 | 0 |
| Total | 507,599 | 444,908 | 796 | 953,302 | 11 | -12,802 | 1,366 |

Table 18. Ontario 2013¹³ Water Use Data Summary

In millions of liters per day

Water Sources: Great Lakes surface water (GLSW), other surface water (OSW) and groundwater (GW)

Table 19. Ontario 2013¹⁴ Intrabasin Diversion Summary

| Sector | Wate | Intrabasin | |
|---------------------|--------------|--------------|-----------|
| | Source | Receiving | Diversion |
| Public Water Supply | Lake Huron | | 159 |
| Public Water Supply | Lake Ontario | | 4 |
| Public Water Supply | | Lake Erie | -151 |
| Other Self Supply | Lake Erie | | 15,978 |
| Other Self Supply | | Lake Ontario | -15,978 |

In millions of liters per day

¹³ 2013 water use data was used for the 2014 report.

¹⁴ 2009 water use data was used for the intrabasin transfers for public supply purposes and 2013 water use data was used for the intrabasin transfer for other supply purposes.

Pennsylvania

The Pennsylvania Lake Erie watershed spans 1,316 square kilometers. The largest land uses in Pennsylvania's portion of the Basin are agriculture and forest.¹⁵ While it is the smallest watershed in the state, it is home to more than 240,000 people with the majority concentrated along the 123 kilometers of Lake Erie coastline.

The total withdrawal amount from the Basin for Pennsylvania was 138 mld. The majority (115 mgd or 83% of the total withdrawal amount) was used for public water supply purposes.

No diversions were reported in 2014. The total consumptive use was 23 mld. The public water supply sector made up the majority (51%) of the total consumptive use amount.

The water use data were provided by the Pennsylvania Department of Environmental Protection (DEP). Reporting compliance varied among water use sectors from 21 percent for the irrigation sector to 100 percent for the self-supply commercial and institutional sector. Pennsylvania DEP did not include estimated water use for the facilities that failed to report their water use to the state.

| Sector | | With | drawals | | Diver | sions | Consumptive |
|---|------|------|---------|-------|------------|------------|-------------|
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 107 | 0 | 7 | 115 | 0 | 0 | 11 |
| Self-Supply Commercial & Institutional | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Self-Supply Irrigation | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Self-Supply Livestock | 0 | 7 | 5 | 12 | 0 | 0 | 10 |
| Self-Supply Industrial | 10 | 0 | 0 | 10 | 0 | 0 | 1 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Self Supply | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 117 | 8 | 12 | 138 | 0 | 0 | 23 |

Table 20. Pennsylvania 2014 Water Use Data Summary

In millions of liters per day

¹⁵ Email communications with David Skellie, Pennsylvania Sea Grant. 2013.

Québec

The majority of Québec's population lives in the Great-Lakes St. Lawrence River watershed. The portion of the St. Lawrence River included in the Great Lakes – St. Lawrence Basin Agreement territory includes the Montreal metropolitan area that represents nearly 50 percent of Québec's population. Some of the tributaries with the greatest flow within that portion are the Outaouais (Ottawa) River, the Richelieu River and the St. François River.

The total withdrawal amount from the Basin for Québec was 4,751 mld - a 1.9 percent decrease from the 2013 withdrawal total of 4,663 mld. The majority (72%) of this amount was used for public water supply purposes at 3,437 mld. The next major water use, industrial sector, made up a quarter of the total withdrawals at 1,228 mld.

The total diversion amount was 12 mld for public supply purposes. The total consumptive use amount was 1,565 mld, representing 33 percent of the total withdrawal amount. The primary water use sectors contributing to the total consumptive use were public supply at 988 mld and industrial at 541 mld.

Starting with 2012 water use data, the province of Québec began its water use data collection program which gathers estimated or metered water use data reported by water users. Being rather new, this program has made progress in identifying and correcting reporting errors, and increasing reporting compliance for all the water use sectors. Québec will begin to collect water use reports from the irrigation (agricultural users) and livestock users in 2016. Therefore, the 2014 data for this sector should be considered as incomplete and unrepresentative of this water use sector in Québec.

| Sector | | Withdr | awals | | Diver | sions | Consumptive |
|---|-------|--------|-------|-------|------------|------------|-------------|
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use |
| Public Water Supply | 2,394 | 843 | 200 | 3,436 | 0 | 12 | 988 |
| Self-Supply Commercial & Institutional | 2 | 14 | 2 | 18 | 0 | 0 | 8 |
| Self-Supply Irrigation | 1 | 9 | 2 | 12 | 0 | 0 | 10 |
| Self-Supply Livestock | 0 | 1 | 0 | 1 | 0 | 0 | 0 |
| Self-Supply Industrial | 537 | 603 | 88 | 1,228 | 0 | 0 | 541 |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Self Supply | 43 | 12 | 1 | 56 | 0 | 0 | 19 |
| Total | 2,977 | 1,481 | 292 | 4,751 | 0 | 12 | 1,565 |

Table 21. Québec 2014 Water Use Data Summary

In millions of gallons per day

Wisconsin

About 25,900 square kilometers of Lakes Michigan and Superior lie within Wisconsin's borders.¹⁶ The state has more than 1,609 kilometers of Great Lakes shoreline and more than 20 percent of the state's land area lies within the Basin where half the population of the state also lives. More than 1.6 million Wisconsin citizens get their drinking water from Lake Michigan or Lake Superior.¹⁷

The total reported water withdrawal amount from the Basin for Wisconsin was 14,989 mld, a five percent decrease from the 2013 water withdrawal total of 15,800 mld. The majority (99%) of the withdrawals came from the Lake Michigan watershed. The primary water use sectors were thermoelectric power production (once-through and recirculated cooling) at 12,835 mld (86% of the total withdrawal amount), public water supply at 1,181 mld, and industrial at 672 mld.

The total reported diversion was 75 mld from the Lake Michigan watershed, mainly for thermoelectric power production (recirculated cooling) purposes. The total consumptive use was 1,348 mld. The primary consumptive uses came from the thermoelectric power (1,057 mld), irrigation (98 mld) and public water supply (117 mld) sectors.

The water use data were provided by the Wisconsin Department of Natural Resources. Reporting compliance varied among water use sectors from 94 percent for the industrial sector to 100 percent for the public supply sector. Data was not estimated for the facilities that did not report water use.

| • | | | | | | | | |
|---|--------|--------|-------|--------|------------|------------|-------------|--|
| Sector | | Withdr | awals | | Diver | sions | Consumptive | |
| Sector | GLSW | OSW | GW | TOTAL | Intrabasin | Interbasin | Use | |
| Public Water Supply | 913 | 81 | 187 | 1,181 | 0 | 25 | 117 | |
| Self-Supply Commercial & Institutional | 5 | 8 | 8 | 22 | 0 | 0 | 3 | |
| Self-Supply Irrigation | 0 | 11 | 130 | 142 | 0 | 0 | 98 | |
| Self-Supply Livestock | 0 | 56 | 56 | 112 | 0 | 0 | 17 | |
| Self-Supply Industrial | 0 | 613 | 59 | 672 | 0 | 0 | 55 | |
| Self-Supply Thermoelectric Power Production (Once-through cooling) | 11,772 | 925 | 0 | 12,697 | 0 | 0 | 1,027 | |
| Self-Supply Thermoelectric Power Production (Recirculated cooling) | 138 | 0 | 0 | 138 | 0 | 49 | 30 | |
| Off-Stream Hydroelectric Power Production | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| In-Stream Hydroelectric Water Use | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other Self Supply | 18 | 4 | 3 | 25 | 0 | 0 | 1 | |
| Total | 12,847 | 1,699 | 443 | 14,989 | 0 | 75 | 1,348 | |

Table 22. Wisconsin 2014 Water Use Data Summary

In millions of liters per day

¹⁶ Wisconsin Sea Grant. 2013. http://seagrant.wisc.edu/Home/AboutUsSection/PressRoom/Details.aspx?PostID=796

¹⁷ Wisconsin Department of Natural Resources. 2013. http://dnr.wi.gov/topic/greatlakes/learn.html

Appendices

Appendix A. Revised protocols for the collection and reporting of water use data

The 2013 regional water use data set is the first to be completed under the new data reporting protocols. The new database and this annual report are expanded and enhanced with the inclusion of new and reformatted water use data and information. Summarized below are the main changes to the database, which are reflected in this report.

- <u>Water use sectors</u> The number of sectors increased from 9 to 10. The Self-Supply Domestic sector is replaced with Self-Supply Commercial and Institutional. The Hydroelectric Power sector is broken down into Off-stream Hydroelectric Power Production and In-stream Hydroelectric Water Use.
- 2. <u>Diversions</u> Three new fields (diversion return flow, net diversion change and diversion return) were added to meet the standards set forth in the new protocols.
- 3. <u>Intrabasin transfers</u> Two additional fields, intrabasin return flow and intrabasin consumptive use, were added to meet the standards set forth in the protocols.
- 4. <u>Two new consumptive use (CU) data fields</u> These fields now more accurately describe how the CU amounts are calculated. The first field documents the percentage of reported consumptive uses that were determined through actual measurement. The second field documents the coefficient or the range of coefficients used to calculate CU. Under the old database regime, a summary chart of the range of CU coefficients was inserted in the annual reports for reference.
- 5. <u>Aggregation</u> The level of aggregation of data was dropped since the protocols do not call for such a field. In its place a "methods" field describes the method used to determine withdrawal amount.

Appendix B. General Definitions from the Compact and Agreement

Basin or Great Lakes-St. Lawrence River Basin means the watershed of the Great Lakes and the St. Lawrence River upstream from Trois-Rivières, Québec.

Consumptive Use means that portion of the water withdrawn or withheld from the basin that is lost or otherwise not returned to the basin due to evaporation, incorporation into products or other processes.

Diversion means a transfer of water from the basin into another watershed, or from the watershed of one of the Great Lakes into that of another by any means of transfer, including but not limited to a pipeline, canal, tunnel, aqueduct, channel, modification of the direction of a water course, a tanker ship, tanker truck or rail tanker but does not apply to water that is used in the basin or a Great Lake watershed to manufacture or produce a product that is then transferred out of the basin or watershed.

Divert has a corresponding meaning.

Withdrawal means the taking of water from surface water or groundwater.

Source Watershed means the watershed from which a withdrawal originates. If water is withdrawn directly from a Great Lake or from the St. Lawrence River, then the Source Watershed shall be considered to be the watershed of that Great Lake or the watershed of the St. Lawrence River, respectively. If water is withdrawn from the watershed of a stream that is a direct tributary to a Great Lake or a direct tributary to the St. Lawrence River, then the Source Watershed of that Great Lake or the watershed shall be considered to be the watershed of that Great Lake or the St. Lawrence River, then the Source Watershed shall be considered to be the watershed of that Great Lake or the watershed of the St. Lawrence River, respectively, with a preference to the direct tributary stream watershed from which it was withdrawn.

Appendix C. Water Use Sector Definitions

Public Water Supply

Water distributed to the public through a physically connected system of treatment, storage and distribution facilities serving a group of largely residential customers that may also serve industrial, commercial and other institutional operators. Water withdrawn directly from the basin and not through such a system shall not be considered to be used for Public Water Supply purposes.

Self-Supply Commercial and Institutional

Commercial uses include water used by motels, hotels, restaurants, office buildings and institutions, both civilian and military. This category also includes water for mobile homes, hospitals, schools, air conditioning and other similar uses not covered under a public supply. In addition, this category includes amusement and recreational water uses such as snowmaking and water slides.

Self-Supply Irrigation

Water artificially applied on lands to assist in the growing of crops and pastures or in the maintenance of recreational lands, such as parks and golf courses.

Self-Supply Livestock

Water used by animals such as horses, cattle, sheep, goats, hogs and poultry. Water used in fish hatchery operations is also included under this category.

Self-Supply Industrial

Industrial water includes water used in the manufacture of metals, chemicals, paper, food and beverage and other products. Mining water use includes water used in the extraction or washing of minerals, for example solids, such as coal and ores, and liquids such as crude petroleum and natural gas. Water used in quarrying and milling is also included in the industrial category. Brine extraction from oil and gas operations is not included. Withdrawals and consumptive uses for industrial and mining purposes (including dewatering operations) recorded under another category (e.g., public supply) will not be recorded here. Once initially reported, water used in a closed cycle (recirculation) will not be reported as a withdrawal. "Make-up water¹⁸" will be reported once upon entering the system. Other situations should be evaluated on a case-by-case basis.

Self-Supply Thermoelectric Power Production (Once-through cooling)

Withdrawals and consumptive uses already recorded under another category (e.g., public supply) will not be reported here.

Self-Supply Thermoelectric Power Production (Recirculated cooling)

Withdrawals and consumptive uses already recorded under another category (e.g., public supply) will not be reported here. Once initially reported, water used in a closed cycle (recirculation) will not be reported as a withdrawal. "Make-up water" will be reported once upon entering the system.

¹⁸ For industrial boiler systems, make-up water is the raw water, softened water or demineralized water required for steam generation. http://www.pdhcenter.com/courses/m165/m165content.pdf

Off-Stream Hydroelectric Power Production

Water removed from a stream channel and used to drive turbines that generate electric power. This category also includes "off-stream use" for pumped-storage systems [e.g., reservoir storage] that return water to the source.

In-Stream Hydroelectric Water Use

This category includes "run of the river" use, which is not considered a water withdrawal or consumptive use. Reporting for this category is voluntary.

Other Self Supply

Water used for purposes not reported in categories 1-9. Examples include, but are not limited to, withdrawals for fish/wildlife, environmental, navigation and water quality purposes. Specifically, water used to maintain levels for navigation, for fish and wildlife habitat creation and enhancement (excluding fish hatchery operations included in category four), for flow augmentation (or diversion), for sanitation, pollution confinement, and other water quality purposes, and agricultural activities (services) other than those directly related to irrigation.

Appendix D. Interim Cumulative Impact Assessment

Executive Summary

This interim cumulative impact assessment, as part of the 2014 Annual Water Use Report, covers the years 2013 and 2014. It is considered interim assessment and is the third in a series of cumulative impact assessments for the Great Lakes and St. Lawrence River basin (Basin).

- 1. *Cumulative Impact Assessment of Withdrawals, Consumptive Uses and Diversions*, published in 2013, covered 2006 to 2010 (2006-2010 Cumulative Impact Assessment).
- 2. Interim Cumulative Impact Assessment of the 2013 Annual Water Use Report, published in 2014, compared the years 2011 and 2012 (2011-2012 Interim Assessment).
- 3. *Interim Cumulative Impact Assessment of the 2014 Annual Water Use Report*, provided in this report (this interim assessment), covers the 2013 and 2014.

These cumulative impact assessments fulfill the requirements of the Great Lakes-St. Lawrence River Basin Sustainable Water Resources Agreement and the companion Great Lakes-St. Lawrence River Basin Water Resources Compact. While the 2006-2010 Cumulative Impact Assessment was the first full assessment covering a five year period, the subsequent, interim cumulative impact assessments have been prepared to track annual water loss to the Basin between full cumulative impact assessments. Water loss is defined as consumptive uses and diversions less return flow. It reflects water not being returned to the source watershed.

The approach used for this interim assessment is similar to that of the 2006-2010 Cumulative Impact Assessment. The analysis focuses on the hydrologic effects of consumptive uses and diversions on water supply and flow at the watershed (i.e., lake basin) scale as well as for the entire Basin.

For comparative purposes, longer data sets for flows, covering the period from 1948 to 2010, were presented to provide a historical context for the assessment. The Basin water budget was used to account for the water flows into and out of the Basin as outlined below.

1. Accounting for Inflows

The inflows included precipitation on the surface of the Great Lakes, surface water runoff to the Great Lakes or the St. Lawrence River, diversions, and connecting channel flows into each of the Great Lakes or the St. Lawrence River, except for Lake Superior which is the headwater to the system.

2. Accounting for Outflows

Outflows included evaporation from the surface of the Great Lakes, diversions from some Lake watersheds, connecting channel flows out of each of the Lakes, and consumptive uses. The St. Lawrence River is the outflow for Lake Ontario and for the entire Basin.

3. Hydrologic Effect Assessment

Although withdrawals are a component of the water budget, the 2006-2010 Cumulative Impact Assessment considered only the hydrologic effect of consumptive uses and diversions. The hydrologic effect is defined as the consumptive uses plus net diversions. Consumptive use is defined as the portion of water withdrawn but not returned due to evaporation, incorporation into products, and other processes.

The following observations were made in the 2006-2010 Cumulative Impact Assessment:

- Diversions and consumptive use were very small relative to inflows. The cumulative hydrologic effect of consumptive uses and diversions were small relative to inflows. While inflows fluctuated from 2006-2010, the cumulative hydrologic effect of consumptive uses and diversions was fairly constant for this time period. The net effect of consumptive uses and diversions was positive for the Basin's water budget. In other words, more water was diverted into the Basin than the total combined amount of water diverted out of the Basin or withdrawn and not returned.
- The uncertainty associated with estimated inflow and outflow data was significantly larger than total consumptive use for the Basin.

It is difficult to assess the cumulative impact of diversions and consumptive uses apart from the natural variability of inflows and outflows of the Great Lakes-St. Lawrence River system. Uncertainty in the Basin water budget components was more than twelve and a half times the total reported consumptive uses in 2008. To illustrate, total runoff to the Basin in 2008 was 259,888 cubic feet per second (cfs). Assuming a 15 percent uncertainty, the amount of calculated runoff may be off by over 38,000 cfs. In comparison, consumptive use in 2008 was only 3,016 cfs. As a result of this, the hydrologic effects of consumptive uses on flows and water levels are difficult to discern relative to uncertainties in the natural inflows and outflows.

The 2006-2010 Cumulative Impact Assessment includes detailed information on the definitions, methodology, assumptions, uncertainty, data sources used as well as specific factors affecting each watershed. Refer to that assessment for clarification of the methods used in this assessment. The full report, *Cumulative Impact Assessment of Withdrawals, Consumptive Uses and Diversions, 2006-2010*, is available at http://glslregionalbody.org/ or http://www.glslcompactcouncil.org/.

The most recent data submitted to the Great Lakes-St. Lawrence Regional Water Use Database indicate a reported increase in incremental water losses¹⁹ to the Basin between 2012 to 2013 of 1,407 cubic feet per second [cfs] (909 mgd). From 2013 to 2014, the total reported water loss for Basin decreased by 722 cfs (467 mgd). A more detailed description of these water losses for 2013 and 2014 are provided in the diversion and consumptive uses section of the 2014 Annual Water Use Report.

For this interim assessment, the following observations are made:

- For 2013 and 2014, more water left the Basin (from consumptive uses and outgoing diversions) than entered the Basin from incoming diversions.
- For 2013 and 2014, diversions and consumptive uses remained very small relative to inflows. For example, diversions and consumptive uses reported for the Lake Superior watershed in 2014 made up 4.21 percent of the average total inflows (from 1948-2010) into the watershed. For the Lake Michigan-Huron watershed in 2014, diversions and consumptive uses made up 1.32 percent of the average total inflows into the watershed over this same period.

Approach

Similar to the approach of pervious assessments, this interim assessment focuses on the hydrologic effects of consumptive uses and diversions on water supply and flow at watershed and Basin scales. These hydrologic effects are presented in the context of watershed and Basin water budgets. The analysis focuses on the consumptive uses and diversions components of the water budget, instead of describing and analyzing all components of the water budget. Annual estimates of inflows and outflows are not provided for the years 2012, 2013 and 2014. Supplemental inflow data presented as a 62-year average (1948-2010)


¹⁹ Incremental water loss is defined as new or increased outgoing diversions plus consumptive uses.

in the 2006-2010 Cumulative Impact Assessment are provided for each watershed and the Basin for comparative purposes.

Following standard scientific procedures, inflows are presented as positive numbers and outflows are presented as negative numbers. This convention is used to help relate different flows to one another and to supply. It is not intended to communicate the effect of these flows on the Basin. All flows are given in cubic feet per second (cfs).

Great Lakes-St. Lawrence River Basin

Figure 1 shows diversions and consumptive uses for the Basin by year for 2012-2014. Net diversions are shown as positive (or incoming) flows into the Basin, mainly due to the Long Lac and Ogoki diversions which divert water from the Hudson Bay watershed into Lake Superior for power generation purposes. These diversions have offset the consumptive uses (displayed as negative numbers in Figure 2) and outgoing diversions.



Diversions Consumptive uses

Figure 1. Diversions and Consumptive Uses for the Great Lakes – St. Lawrence Basin




Figure 2. Net Diversions and Consumptive Uses for the Great Lakes – St. Lawrence River Basin.

The numbers in Table A below (presented graphically in Figure 2), indicate for the Basin the cumulative hydrologic effect of consumptive uses and diversions (annual averages) are small relative to inflows (runoff plus precipitation). In general, the cumulative hydrologic effect of consumptive uses and diversions has fluctuated for 2012-2014. Between the years 2012 to 2013 consumptive uses and diversions increased by 1,407 cfs (96 cfs to -1,311 cfs). From 2013 to 2014, consumptive uses and diversions for Basin decreased by 741 cfs (from -1,311 cfs to -570 cfs).

For 2012, the net effect of consumptive uses and diversions was positive for the Basin. In other words, more water was diverted into the Basin than the total combined amount of water diverted out of the Basin or withdrawn and not returned. In 2013 and 2014 the net effect of consumptive uses and diversions was negative for the Basin. More water was consumptively used or diverted out of the Basin than the total combined amount of water diverted into the Basin.

| Year | Runoff + Precipitation | Consumptive Uses + Diversions |
|------|------------------------|-------------------------------|
| 2012 | 434,161* | 96 |
| 2013 | 434,161* | -1,311 |
| 2014 | 434,161* | -570 |

Table A. Water budget values in cubic feet per second for the Great Lakes-St. Lawrence River Basin, 2012-2014. *62-year flow average

Lake Superior Watershed

The data in Figure 3 and Table B summarize the hydrologic effects of consumptive uses and diversions for the Lake Superior watershed water budget. For purposes of comparison, Figure 3 depicts all components of the water budget (e.g., runoff, precipitation, evaporation and the flow of the St. Marys River) using the annual average for the years 1948-2010.




Figure 3. Water budget average flows for Lake Superior, comparing 2012-2014 a historical 62-year period (1948-2010). (in cfs)

| Water Budget Component | 1948-2010 62-year Flow | 2012 | 2013 | 2014 |
|---------------------------|---------------------------|-------|-------|-------|
| Diversions | 5,950 | 5,863 | 4,651 | 5,220 |
| Consumptive Uses | -117 | -70 | -73 | -71 |

Table B. Diversions and consumptive uses for Lake Superior, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

Consumptive Uses: Consumptive uses in the Lake Superior watershed have remained relatively constant over the years from 2012 to 2014.

Diversions: Diversions, mainly comprised of the Long Lac and Ogoki Diversions, fluctuate with the weather conditions of the watersheds. When conditions in the Long Lac and Nipigon (downstream of Ogoki) watersheds are wet, the diversions are often reduced, and water that otherwise would have been diverted into Lake Superior is instead directed through natural outlets that flow toward Hudson Bay. Conversely, when conditions are dry in the downstream watersheds, the diversion flow may be higher.

As illustrated in Table C below, for the Lake Superior watershed the hydrologic effect of consumptive uses and diversions (annual averages) are small relative to inflows. The estimated net volume of consumptive uses and diversions decreased by 34 percent from 2012 to 2013, and then increased from 2013 to 2014 by 12 percent.

| Year | Total Inflow | Consumptive Uses +Diversions | Consumptive Uses + Diversions (as a percentage of total inflow) |
|--------------|-----------------|------------------------------|--|
| 62-year avg. | 122,218* | 5,833 | 4.77% |
| 2012 | 122,218* | 6,957 | 5.69% |
| 2013 | 122,218* | 4,578 | 3.75% |
| 2014 | 122,218* | 5,148 | 4.21% |

Table C. Water budget values in cubic feet per second for Lake Superior, 62-year average for 1948-2010, 2012-2014. (in cfs)

*62-year flow average

Lakes Michigan-Huron Watershed

The data in Figure 4 and Table D summarize the hydrologic effects of the consumptive use and diversion components of the Lakes Michigan-Huron watershed water budget. For purposes of comparison, Figure 4 depicts the all components of the water budget using the annual average for the years1948-2010.




Figure 4. Water budget average flows for Lakes Michigan-Huron, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

| Water Budget Component | 62-year Flow | 2012 | 2013 | 2014 |
|------------------------|--------------|--------|--------|--------|
| Diversions | -3,171 | -1,894 | -1,883 | -1,885 |
| Consumptive Uses | -1,166 | -1,779 | -1,701 | -1,839 |

Table D. Diversions and consumptive uses for Lakes Michigan-Huron, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

Consumptive Uses: Consumptives use flows fluctuated from 2012 to 2014.

Diversions: Diversions remained relatively constant from 2012-2014. Diversions, mainly comprised of the Illinois Diversion which diverts water from Lake Michigan to the Chicago Sanitary and Ship Canal and the Illinois and Des Plaines Rivers, contributed to the flows out of the watershed.

As illustrated in Table E, for the Lakes Michigan-Huron watershed the hydrologic effect of consumptive uses and diversions (annual averages) were small relative to inflows (about 1.3% of the 62-year average for inflows into the watershed). The estimated net volume of diversions and consumptive uses decreased from 2012 to 2013 by 13 percent and remained relatively the same between 2013 and 2014.

| Year | Total Inflow | Consumptive Uses +Diversions | Consumptive Uses + Diversions (as a percentage of total inflow) |
|--------------|-----------------|------------------------------|--|
| 62-year avg. | 282,054* | -4,337 | 1.53% |
| 2012 | 282,054* | -4,413 | 1.56% |
| 2013 | 282,054* | -3,799 | 1.35% |
| 2014 | 282,054* | -3,723 | 1.32% |

 Table E. Water budget values in cubic feet per second for Lakes Michigan-Huron, 62-year average for 1948-2010, 2012-2014. (in cfs)

*62-year flow average

Lake Erie Watershed

The data in Table F and used in Figure 5 summarize the hydrologic effect of the consumptive use and diversion components of the Lake Erie watershed water budget. For purposes of comparison, Figure 5 depicts all components of the water budget using the annual average for the years 1948-2010. In 2013 and 2014, these diversion and consumptive use flows were smaller than the 62-year average flows and the 2012 values.




Figure 5. Water budget average flows for Lake Erie, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

| Water Budget Component | 62-year Flow | 2012 | 2013 | 2014 |
|------------------------|--------------|--------|--------|--------|
| Diversions | -7,851 | -8,017 | -7,645 | -6,164 |
| Consumptive Uses | -763 | -736 | -689 | -682 |

Table F. Diversions and consumptive uses for Lake Erie, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

Consumptive Uses: Consumptive uses have decreased from 2012 to 2014. The 2014 consumptive use amount of 682 cfs is about 10 percent less than the annual average for the years 1948-2010.

Diversions: Diversions²⁰ have contributed to the overall flows out of the watershed. Similar to the decreasing trend in consumptive uses, diversions have declined from 8,017 cfs in 2012 to 6,164 cfs in 2014.

As illustrated in Table G, for the Lake Erie watershed the hydrologic effect of consumptive uses and diversions (annual averages) was small relative to inflows. The estimated net volume of consumptive uses and diversions decreased significantly from 2012 to 2014.

| Year | Total Inflow | Consumptive Uses +Diversions | Consumptive Uses + Diversions (as a percentage of total inflow) |
|--------------|--------------|---------------------------------|--|
| 62-year avg. | 244,739* | -8,614 | 3.51% |
| 2012 | 244,739* | -8,753 | 3.58% |
| 2013 | 244,739* | -8,333 | 3.41% |
| 2014 | 244,739* | -6,846 | 2.79% |

Table G. Water budget values in cubic feet per second for Lake Erie, 62-year average for 1948-2010, 2012-2014. (in cfs)

*62-year flow average

Lake Ontario Watershed

The data in Table H and used in Figure 6 summarize the hydrologic effects of the consumptive use and diversion components of the Lake Ontario watershed water budget. For purposes of comparison, Figure 6 depicts all components of the water budget using the annual average for the years 1948-2010.




Figure 6. Water budget average flows for Lake Ontario, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

²⁰ Diversion data for the Lake Erie watershed include an intrabasin diversion at Welland Canal.

| Water Budget Component | 62-year Flow | 2012 | 2013 | 2014 |
|------------------------|--------------|-------|-------|-------|
| Diversions | 7,851 | 7,942 | 7,575 | 6,465 |
| Consumptive Uses | -561 | -375 | -567 | -576 |

Table H. Diversions and consumptive uses for Lake Ontario, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

Consumptive Uses: These consumptive use flows appear to have increased from the 2012 to 2014 reported amounts.

Diversions: There was a notable decrease in diversion flows from 7,942 cfs in 2012 to 6,465 cfs in 2014.

As illustrated in Table I, for the Lake Ontario watershed the cumulative hydrologic effect of consumptive uses and diversions (annual averages) were small relative to inflows. The estimated net volume of diversions and consumptive uses decreased by 7 percent from 2012 to 2013 and remained relatively constant from 2013 to 2014.

| Year | Total Inflow | Consumptive Uses +Diversions | Consumptive Uses + Diversions (as a percentage of total inflow) |
|--------------|-----------------|------------------------------|--|
| 62-year avg. | 269,041* | 7,290 | 2.71% |
| 2012 | 269,041* | 7,567 | 2.81% |
| 2013 | 269,041* | 7,009 | 2.61% |
| 2014 | 269,041* | 7,041 | 2.61% |

 Table I. Water budget values in cubic feet per second for Lake Ontario, 62-year average for 1948-2010, 2012-2014. (in cfs)

 ***2
 #**2

*62-year flow average

St. Lawrence River Watershed

The water budget for the St. Lawrence River watershed is different than those for the Lakes. Inflow consists of the St. Lawrence River flow measured at Cornwall, Ontario. Outflow mainly consists of the river's flow modeled at Trois Rivières, Québec and consumptive uses throughout the watershed.

As illustrated in Table K and Figure 7, for the St. Lawrence River watershed the hydrologic effect of consumptive use and diversions was small relative to inflows. From 2012 to 2013, the net estimated volume of consumptive uses and diversion increased by 21 percent (from 632 cfs to 765 cfs), then decreased to 694 cfs in 2014^{21} .

²¹ The implementation of the 2009 Water Use Data Protocols and the corresponding jurisdictional data collection and reporting programs began with the collection of 2012 water use data. Therefore, the increase of the net estimated volume of consumptive uses and diversions between 2012 and 2014 may reflect the improvements in data collection.




Table 7. Water budget values in cubic feet per second for the St. Lawrence River comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

| Water Budget Component | 62-year Flow | 2012 | 2013 | 2014 |
|------------------------|--------------|------|------|------|
| Diversions | N/A | -7 | -9 | -9 |
| Consumptive Uses | -313 | -625 | -756 | -685 |

Table J. Diversions and consumptive uses for St. Lawrence River, comparing 2012-2014 to a historical 62-year period (1948-2010). (in cfs)

Consumptive Uses: As summarized in Table J, consumptive uses have fluctuated, increasing from 625 cfs in 2012 to 756 cfs in 2013, and then decreasing to 685 cfs in 2014.

Diversions: Diversions have made up about 1 percent of the total annual water loss from the St. Lawrence River watershed. Starting in 2012, both Québec and New York reported diversions totaling 7 cfs in 2012, 9 cfs in 2013 and 9 cfs in 2014 for public supply purposes. For previous years (1948-2010), diversions have not been reported.

As illustrated in Table K, for the St. Lawrence River watershed the cumulative hydrologic effect of consumptive uses and diversions (annual averages) were small relative to inflows.

| Year | Total Inflow | Consumptive uses +Diversions | Consumptive Uses + Diversions (as a percentage of total inflow) |
|--------------|-----------------|------------------------------|--|
| 62-year avg. | 256,797* | -313** | 0.12% |
| 2012 | 256,797* | -632 | 0.25% |
| 2013 | 256,797* | -765 | 0.30% |
| 2014 | 256,797* | -694 | 0.27% |

Table K. Water budget values in cubic feet per second for the St. Lawrence River, 62-year average for 1948-2010, 2012-2014. (in cfs)

```
*62-year flow average
```

**This figure only is consumptive use only. No diversions were reported.